

publ. 1995

TRICHOSCAPA KARAWAJEWI AND ITS SYNONYMS
(HYMENOPTERA: FORMICIDAE)

BY WILLIAM L. BROWN, JR.

Department of Entomology
Cornell University
Ithaca, NY 14853

A recent loan from the Zoological Institute, Ukrainian Academy of Sciences, Kiev, Ukraine, through the kindness of Dr. Alexander G. Radchenko, has allowed me for the first time to examine the type of *Strumigenys emeryi* Karawajew, which proves to be an example of the same species I later described as *Smithistruma dubia*. Since *S. emeryi* was preoccupied in *Strumigenys*, I had already given it the replacement name *Strumigenys (Trichoscapa) karawajewi*. The necessary synonymy follows.

***Trichoscapa karawajewi* (new combination)**

Strumigenys (Cephaloxys) emeryi Karawajew, 1935, *Treubia* 15:106-108, fig. 25, queen, type locality Siak, Sumatra. Preoccupied.

Strumigenys (Trichoscapa) karawajewi Brown, 1948, *Entomological News*, Philadelphia, 59:44, replacement name.

Smithistruma (Smithistruma) dubia Brown, 1953, *American Midland Naturalist*, 50:122, worker, queen, type locality East Coast of Peleliu I., Palau Group. New synonym.

In the type from Kiev, the clypeus is broader than long. In addition, the larger hairs on the head are as arranged in other samples of this species, and are not restricted to two at cephalic midlength as depicted by Karawajew. Although I described *S. dubia* from atolls in the South Pacific, while *S. karawajewi* came from Sumatra, we now have samples of this species from numerous sites in New Guinea and the Solomons, Sulawesi, Luzon, Sabah, Manus, Truk, and Ponape, as well as Fiji. Because the limits of the short-mandibulate dacetite dacetine genera have grown very unclear as